arXiv:1207.1630v4 [q-fin.CP] 31 Jul 2012

Local Lévy Models and their Volatility Smile

Matthew Lorig *

This version: August 1, 2012

Abstract

We propose a class of equity models whose volatility, Lévy measure, and killing rate all have local stochastic state-dependence. In this framework we find a closed form solution for the price of any European-style option. Additionally, for a certain sub-class of models, we find an exact expression for the induced implied volatility smile. To illustrate our framework, we perform specific computations for hybrid CEV/Lévy model (which we christen as the "CÉV" model).

Keywords: CEV, Lévy, Local Volatility, Implied Volatility, Default

1 Introduction

A *local volatility* model is a stochastic volatility model in which the volatility σ_t of an asset X is a function of the present level of X. That is, $\sigma_t = \sigma(X_t)$. Among local stochastic volatility models, perhaps the most well-known is the constant elasticity of variance (CEV) model of Cox (1975). An extension of the CEV model to defaultable assets (the Jump-to-Default CEV or JDCEV model) is derived in Carr and Linetsky (2006). One advantage of these models is that they admit closed-form pricing formulas (written as infinite series of special functions) for European options. However, other than a possible jump to default, neither of these models (or any other local stochastic volatility model) allows the underlying asset to experience jumps (though jumps can be inherited through Lévy subordination, as described in Mendoza-Arriaga, Carr, and Linetsky (2010)).

Exponential Lévy models, on the other hand, *do* allow the underlying price process to experience jumps. Furthermore, in an exponential Lévy setting, European option-prices can be computed quickly as generalized Fourier transforms (see Lewis (2001); Lipton (2002)). However, exponential Lévy models are spatially homogeneous; neither the volatility nor the jump-intensity not have any local dependence. Thus, exponential Lévy models do not exhibit volatility clustering or the leverage effect.

---

*ORFE Department, Princeton University, Princeton, USA. Work partially supported by NSF grant DMS-0739195

In this paper, we introduce a class of *local Lévy models*. Such models allow for the volatility, Lévy measure, and killing intensity of an asset to have stochastic local dependence. Within this framework, we find a closed form solution for the price of any European style option and, for certain sub-class of models, we find a closed form solution for the induced implied volatility smile.

The rest of this paper proceeds as follows: in section 2 we present our class of models and our assumptions. In section 3 we derive a formula for the price of any European option. In section 4 we provide a formula for the implied volatility smile induced by a certain sub-class of models within our framework. And in section 5 we perform specific computations for a CEV-like local Lévy model. Numerical examples are provided at the conclusion of the text. An appendix with some mathematical background is also provided. Some concluding remarks are found in section 6.

2 Model and assumptions

We assume a frictionless market, no arbitrage and take an equivalent martingale measure $\mathbb{P}$ chosen by the market on a complete filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t, t \geq 0\}, \mathbb{P})$. The filtration $\{\mathcal{F}_t, t \geq 0\}$ represents the history of the market. All processes defined below live on this space. For simplicity we assume zero interest rates and no dividends. Thus, all traded assets are martingales. We consider a risky asset $X = \exp(Y)$ with lifetime ζ whose dynamics are given by

$$\begin{aligned} dY_t &= \alpha(Y_t)dt + \Big(a_0^2 + \varepsilon\, a_1^2 \eta(Y_t)\Big)^{1/2} dW_t + \int_{\mathbb{R}} z\, d\widetilde{N}_t(Y_{t-}, dz), & Y_0 = y, \\ \alpha(Y_t) &= \Big(c_0 + \varepsilon\, c_1 \eta(Y_t)\Big) - \frac{1}{2}\Big(a_0^2 + \varepsilon\, a_1^2 \eta(Y_t)\Big) - \int_{\mathbb{R}} \nu(Y_{t-}, dz)\Big(e^z - 1 - z\Big), \\ \zeta &= \inf\left\{ t > 0 : \int_0^t \Big(c_0 + \varepsilon\, c_1 \eta(Y_s)\Big)\, ds = \mathcal{E} \right\}. \end{aligned}$$

Here, $(a_0, a_1, c_0, c_1, \varepsilon)$ are non-negative constants, $\eta \geq 0$ is a function in $C_0^\infty(\mathbb{R})$ (the set of smooth functions on $\mathbb{R}$ with compact support), W is a Brownian motion, $\mathcal{E}$ is an independent exponentially distributed random variable with parameter one, and $d\widetilde{N}_t(Y_{t-}, dz)$ is a compensated Poisson random measure

$$d\widetilde{N}_t(Y_{t-}, dz) = dN_t(Y_{t-}, dz) - \nu(Y_{t-}, dz), \qquad \nu(Y_{t-}, dz) = \nu_0(dz) + \varepsilon\, \eta(Y_{t-})\nu_1(dz).$$

We assume that the locally-dependent Lévy measure $\nu(y, dz)$ satisfies

$$\int_{\mathbb{R}} \min(1, z^2)\nu(y, dz) < \infty, \qquad \int_{|z|\geq 1} e^z \nu(y, dz) < \infty, \qquad \int_{|z|\geq 1} |z|\nu(y, dz) < \infty,$$

for all $y \in \mathbb{R}$. The first integrability condition must be satisfied by all Lévy measures. The second integrability condition is needed to ensure $\mathbb{E}\, X_T < \infty$ for all $T \in \mathbb{R}^+$. The last integrability condition can be removed.

However, keeping this condition vastly simplifies the formulas that follow, as it will allow us to replace the indicator function that usually appears in Lévy integrals with the constant one (i.e. $\mathbb{I}_{\{|z|<1\}} \to 1$). We list the main features of our class of models below.

- The process Y experiences local stochastic volatility with volatility function $\sigma(y) = \left(a_0^2 + \varepsilon\, a_1^2 \eta(y)\right)^{1/2}$.
- Jumps in Y of size dz arrive as a Poisson process with an instantaneous state-dependent intensity of $\nu(Y_{t-}, dz)$. The state-dependent Lévy measure has the decomposition $\nu(y, dz) = \nu_0(dz) + \varepsilon\, \eta(y)\nu_1(dz)$, where ν_0 and ν_1 are Lévy measures. Note that both the jump intensity and jump distribution can change depending on the value of y.
- The process Y is killed with a state-dependent intensity of $k(Y_t) = (c_0 + \varepsilon\, c_1 \eta(Y_t))$. When Y is killed it is sent to a cemetery state Δ, from which it cannot return.
- The drift function $\alpha(y)$ is fixed by the volatility and killing rate so that $X = \exp(Y)$ is a martingale.
- Both negative infinity and infinity are natural boundaries for the process Y according to Feller's boundary classification for one-dimensional diffusions (see Borodin and Salminen (2002) pp. 14-15). Thus, the state-space of Y is $\mathbb{R} \cup \{\Delta\}$.

3 Option Pricing

We wish to find the time-zero value $u^\varepsilon(t, y)$ of a European-style option with payoff $h(Y_t)\mathbb{I}_{\{\zeta>t\}}$ at time $t > 0$. Using risk-neutral pricing we have

$$u^\varepsilon(t, y) = \mathbb{E}_y\, h(Y_t)\mathbb{I}_{\{\zeta>t\}},$$

where the notation $\mathbb{E}_y$ indicates expectation starting from y. The function $u^\varepsilon(t, y)$ satisfies the Kolmogorov backward equation

$$(-\partial_t + \mathcal{A}^\varepsilon)\, u^\varepsilon = 0, \qquad\qquad u^\varepsilon(0, y) = h(y), \tag{1}$$

where $\mathcal{A}^\varepsilon$ is the generator of the process Y. The domain of $\mathcal{A}^\varepsilon$ is defined as the set of f for which the limit $\lim_{t\to0} \frac{1}{t}\left(\mathbb{E}_y f(Y_t)\mathbb{I}_{\{\zeta>t\}} - f(y)\mathbb{I}_{\{\zeta>0\}}\right)$ exists in the strong sense. For any $f \in C_0^\infty(\mathbb{R})$ the generator $\mathcal{A}^\varepsilon$ has the explicit representation

$$\begin{aligned}
&\mathcal{A}^\varepsilon = \mathcal{A}_0 + \varepsilon\, \eta\, \mathcal{A}_1, \\
&\mathcal{A}_i = \frac{1}{2} a_i^2 \left(\partial^2 - \partial\right) + c_i\,(\partial - 1) - \int_{\mathbb{R}} \nu_i(dz)\Big(e^z - 1 - z\Big)\partial + \int_{\mathbb{R}} \nu_i(dz)\Big(e^{z\partial} - 1 - z\partial\Big), \quad i \in \{1, 2\}, \\
&\mathrm{dom}(\mathcal{A}_i) = C_0^\infty(\mathbb{R}).
\end{aligned}$$

The symbol ∂ (without a subscript t) indicates a derivative taken with respect to y. Note that the operator ∂ is the generator of the translations: $e^{z\partial}f(y) = f(y+z)$.

Remark 1. Let $\mathcal{H}$ be the Hilbert space $L^2(\mathbb{R}, dy)$ with inner product $(f,g) := \int_{\mathbb{R}} \overline{f}(y)\, g(y)\, dy$ and norm $\|f\| = \sqrt{(f,f)}$. Note that the operators $\{\mathcal{A}_i, i = 1,2\}$ are normal operators in $\mathcal{H}$ and satisfy the following (improper) eigenvalue equations (neither $\mathcal{A}_i$ have any proper eigenvalues)

$$\mathcal{A}_0 \psi_\lambda = \phi_\lambda \psi_\lambda, \qquad \mathcal{A}_1 \psi_\lambda = \chi_\lambda \psi_\lambda, \qquad \psi_\lambda = \frac{1}{\sqrt{2\pi}} e^{i\lambda y},$$

where

$$\begin{aligned} \phi_\lambda &= \frac{1}{2} a_0^2 \left(-\lambda^2 - i\lambda\right) + c_0(i\lambda - 1) - \int_{\mathbb{R}} \nu_0(dz) \Big(e^z - 1 - z\Big) i\lambda + \int_{\mathbb{R}} \nu_0(dz) \Big(e^{i\lambda z} - 1 - i\lambda z\Big), \\ \chi_\lambda &= \frac{1}{2} a_1^2 \left(-\lambda^2 - i\lambda\right) + c_1(i\lambda - 1) - \int_{\mathbb{R}} \nu_1(dz) \Big(e^z - 1 - z\Big) i\lambda + \int_{\mathbb{R}} \nu_1(dz) \Big(e^{i\lambda z} - 1 - i\lambda z\Big). \end{aligned}$$

The eigenfunctions satisfy $(\psi_\lambda, \psi_\mu) = \delta(\lambda - \mu)$. Note that Borel-measurable functions of normal operators (e.g., $g(\mathcal{A}_0)$) are well-defined, as explained in Appendix A.

We seek a solution to Cauchy problem (1) of the form

$$u^\varepsilon = \sum_{n=0}^{\infty} \varepsilon^n \, u_n. \tag{2}$$

Justification for this expansion will be given in Theorem 2. Inserting the expansion (2) into Cauchy problem (1) and collecting terms of like powers of ε we find

$$\begin{aligned} &\mathcal{O}(1): && (-\partial_t + \mathcal{A}_0) u_0 = 0, && u_0(0,y) = h(y), \\ &\mathcal{O}(\varepsilon^n): && (-\partial_t + \mathcal{A}_0) u_n = -\eta \mathcal{A}_1 u_{n-1}, && u_n(0,y) = 0. \end{aligned}$$

The solution to the above equations is

$$\begin{aligned} &\mathcal{O}(1): && u_0(t,y) = e^{t\mathcal{A}_0} h(y), \\ &\mathcal{O}(\varepsilon^n): && u_n(t,y) = \int_0^t ds\, e^{(t-s)\mathcal{A}_0} \eta(y) \mathcal{A}_1 u_{n-1}(s,y). \end{aligned}$$

Next, using the equation (11) from appendix A we obtain

$$\begin{aligned} &\mathcal{O}(1): && u_0(t,y) = \int_{\mathbb{R}} d\lambda\, e^{t\phi_\lambda} (\psi_\lambda, h) \psi_\lambda(y), \\ &\mathcal{O}(\varepsilon^n): && u_n(t,y) = \int_0^t \int_{\mathbb{R}} ds\, d\mu\, e^{(t-s)\phi_\mu} \left(\psi_\mu, \eta \mathcal{A}_1 u_{n-1}(s,\cdot)\right) \psi_\mu(y), \end{aligned}$$

After a bit of algebra, we find an explicit representation for $u_n(t,y)$

$$u_n(t,y) = \underbrace{\int \cdots \int}_{n+1} \left(\prod_{k=0}^{n} d\lambda_k \right) \left(\sum_{k=0}^{n} \frac{e^{t\phi_{\lambda_k}}}{\prod_{j\neq k}^{n} (\phi_{\lambda_k} - \phi_{\lambda_j})} \right) \left(\prod_{k=0}^{n-1} \chi_{\lambda_k} \left(\psi_{\lambda_{k+1}}, \eta\, \psi_{\lambda_k} \right) \right) (\psi_{\lambda_0}, h)\, \psi_{\lambda_n}. \tag{3}$$

We have now obtained a formal expansion (2)-(3) for the price of a European option. The following theorem provides conditions under which the expansion is guaranteed to be valid.

Theorem 2 (Option Price)**.** *Suppose ε satisfies*

$$\varepsilon^2 \leq \inf_{\lambda \in \mathbb{R}} \frac{a^2 + b^2 |\phi_\lambda|^2}{\|\eta\|^2 \cdot |\chi_\lambda|^2}, \tag{4}$$

for some $a \geq 0$ and $b \leq 1$. Then the option price $u^\varepsilon(t,y)$ is given by infinite series (2) *where the $\{u_n\}$ are given by* (3)*.*

Proof. See Appendix B. □

Remark 3. Not that for certain ϕ_λ and χ_λ the bound (4) may be infinite (by choosing a large enough), in which case expansion (2) is valid for all $\varepsilon < \infty$.

4 Implied Volatility

Assumption 4. In this section only we assume $\nu_0 \equiv 0$ and $c_0 = 0$. Thus

$$\phi_\lambda = \frac{1}{2} a_0^2 \left(-\lambda^2 - i\lambda \right).$$

Note that, with this assumption, if $\varepsilon = 0$, then $X = \exp(Y)$ is a geometric Brownian motion. This will be key for our implied volatility analysis.

We fix (t,y) and a call option payoff $h(y) = (e^y - e^k)^+$. Note that

$$(\psi_\lambda, h) = \frac{-e^{k - ik\lambda}}{\sqrt{2\pi}\,(i\lambda + \lambda^2)}, \qquad \mathrm{Im}(\lambda) < -1.$$

The following definitions will be useful:

Definition 5. The *Black-Scholes Price* $u^{BS} : \mathbb{R}^+ \to \mathbb{R}^+$ is defined as

$$u^{BS}(\sigma) := \int d\lambda\, e^{t\phi_\lambda^{BS}(\sigma)} (\psi_\lambda, h) \psi_\lambda, \qquad \phi_\lambda^{BS}(\sigma) = \frac{1}{2}\sigma^2(-\lambda^2 - i\lambda).$$

Definition 6. The *Implied Volatility* is defined implicitly as the unique number $\sigma^\varepsilon \in \mathbb{R}^+$ such that

$$u^{BS}(\sigma^\varepsilon) = u^\varepsilon, \tag{5}$$

where u^ε is as given in Theorem 2.

Remark 7. Note that $u_0 = u^{BS}(a_0)$. As shown in Lorig (2012), when u^ε can be expanded as a power series whose first term corresponds to u^{BS}, one can obtain the exact implied volatility surface corresponding to u^ε.

Remark 8. For $0 < t < \infty$ the existence and uniqueness of the implied volatility σ^ε can be deduced by using the general arbitrage bounds for call prices and the monotonicity of u^{BS}.

Remark 9. Note that u^{BS} is an invertible analytic function that satisfies $\partial_\rho u^{BS}(\sigma) > 0$ for all $\sigma > 0$. By the Lagrange inversion theorem, the inverse $[u^{BS}]^{-1}$ of such a function is also analytic.

Clearly, u^ε is an analytic function of ε (we derived its power series expansion). It is a useful fact that the composition of two analytic functions is also analytic (see Brown and Churchill (1996), section 24, p. 74). Thus, in light of Remark 9, we deduce that $\sigma^\varepsilon = [u^{BS}]^{-1}(u^\varepsilon)$ is an analytic function and therefore has a power series expansion in ε. We write this expansion as follows

$$\sigma^\varepsilon = \sigma_0 + \delta^\varepsilon, \qquad \delta^\varepsilon = \sum_{k=1}^{\infty} \varepsilon^k \sigma_k. \tag{6}$$

Taylor expanding u^{BS} about the point σ_0 we have

$$\begin{aligned}
u^{BS}(\sigma^\varepsilon) &= u^{BS}(\sigma_0 + \delta^\varepsilon) \\
&= \sum_{n=0}^{\infty} \frac{1}{n!} (\delta^\varepsilon \partial_\sigma)^n u^{BS}(\sigma_0) \\
&= u^{BS}(\sigma_0) + \sum_{n=1}^{\infty} \frac{1}{n!} \left(\sum_{k=1}^{\infty} \varepsilon^k \sigma_k \right)^n \partial_\sigma^n u^{BS}(\sigma_0) \\
&= u^{BS}(\sigma_0) + \sum_{n=1}^{\infty} \frac{1}{n!} \left[\sum_{k=1}^{\infty} \left(\sum_{j_1+\cdots+j_n=k} \prod_{i=1}^{n} \sigma_{j_i} \right) \varepsilon^k \right] \partial_\sigma^n u^{BS}(\sigma_0) \\
&= u^{BS}(\sigma_0) + \sum_{k=1}^{\infty} \varepsilon^k \left[\sum_{n=1}^{\infty} \frac{1}{n!} \left(\sum_{j_1+\cdots+j_n=k} \prod_{i=1}^{n} \sigma_{j_i} \right) \partial_\sigma^n \right] u^{BS}(\sigma_0) \\
&= u^{BS}(\sigma_0) + \sum_{k=1}^{\infty} \varepsilon^k \left[\sigma_k \partial_\sigma + \sum_{n=2}^{\infty} \frac{1}{n!} \left(\sum_{j_1+\cdots+j_n=k} \prod_{i=1}^{n} \sigma_{j_i} \right) \partial_\sigma^n \right] u^{BS}(\sigma_0).
\end{aligned} \tag{7}$$

Now, we insert expansions (2) and (7) into (5) and collect terms of like order in ε

$$\begin{aligned}
&\mathcal{O}(1): && u_0 = u^{BS}(\sigma_0), \\
&\mathcal{O}(\varepsilon^k): && u_k = \sigma_k \partial_\sigma u^{BS}(\sigma_0) + \sum_{n=2}^{\infty} \frac{1}{n!} \left(\sum_{j_1+\cdots+j_n=k} \prod_{i=1}^{n} \sigma_{j_i} \right) \partial_\sigma^n u^{BS}(\sigma_0), \qquad k \geq 1.
\end{aligned}$$

Solving the above equations for $\{\sigma_k\}_{k=0}^{\infty}$ we find

$$\begin{aligned} \mathcal{O}(1): \qquad & \sigma_0 = a_0, \\ \mathcal{O}(\varepsilon^k): \qquad & \sigma_k = \frac{1}{\partial_\sigma u^{BS}(\sigma_0)} \left(u_k - \sum_{n=2}^{\infty} \frac{1}{n!} \left(\sum_{j_1+\cdots+j_n=k} \prod_{i=1}^{n} \sigma_{j_i} \right) \partial_\sigma^n u^{BS}(\sigma_0) \right), \qquad k \geq 1. \end{aligned} \tag{8}$$

Remark 10. The right hand side of (8) involves only σ_j for $j \leq k-1$. Thus, the $\{\sigma_k\}_{k=1}^{\infty}$ can be found recursively.

Remark 11. Note that $\partial_\sigma^n u^{BS}(\sigma)$ is easily computed using

$$\partial_\sigma^n u^{BS}(\sigma) = \int d\lambda \left(\partial_\sigma^n e^{t\phi_\lambda^{BS}(\sigma)} \right) (\psi_\lambda, h) \psi_\lambda.$$

Explicitly, up to $\mathcal{O}(\varepsilon^4)$ we have

$$\begin{aligned} \mathcal{O}(\varepsilon): \qquad & \sigma_1 = \frac{u_1}{\partial_\sigma u_0}, \\ \mathcal{O}(\varepsilon^2): \qquad & \sigma_2 = \frac{u_2 - \frac{1}{2!}\sigma_1^2 \partial_\sigma^2 u_0}{\partial_\sigma u_0}, \\ \mathcal{O}(\varepsilon^3): \qquad & \sigma_3 = \frac{u_3 - (\sigma_2\sigma_1\partial_\sigma^2 + \frac{1}{3!}\sigma_1^3\partial_\sigma^3)u_0}{\partial_\sigma u_0}, \\ \mathcal{O}(\varepsilon^4): \qquad & \sigma_4 = \frac{u_4 - (\sigma_3\sigma_1\partial_\sigma^2 + \frac{1}{2}\sigma_2^2\partial_\sigma^2 + \frac{1}{2}\sigma_2\sigma_1^2\partial_\sigma^3 + \frac{1}{24}\sigma_1^4\partial_\sigma^4)u_0}{\partial_\sigma u_0}. \end{aligned}$$

We summarize our implied volatility result in the following theorem:

Theorem 12 (Implied Volatility)**.** *The implied volatility* σ^ε *defined in* (5) *is given explicitly by* (6) *where* $\sigma_0 = \sigma$ *and* $\{\sigma_k\}_{k=1}^{\infty}$ *are given by* (8)*.*

Remark 13. Everything we have done so far is exact. The accuracy of the implied volatility expansion (6) is limited only by the number of terms one wishes to compute.

5 Example: CÉV

The constant elasticity of variance (CEV) model of Cox (1975) improves upon the Black-Scholes model by allowing the volatility to depend on the present level of the underlying through a local volatility function (in the variable $y = \log x$) of the form $\sigma(y) = \sqrt{\varepsilon}\, a_1 e^{\beta y/2}$. This model has enjoyed wide success because (i) it allows for closed-form solutions for European option prices and (ii) when $\beta < 0$, the local volatility function *increases* as $y \to -\infty$ ($x \to 0$), which is consistent with the leverage effect and results in a negative implied volatility skew. Still, the CEV model has some shortcomings. First, volatility drops to zero as $y \to \infty$ ($x \to \infty$). Second, the model does not allow the underlying to jump.

We can correct both of these shortcomings in our framework by allowing $\eta(y) = e_\beta(y) := e^{\beta y}$. In this setting, the dynamics of Y become

$$\begin{aligned} dY_t &= \alpha(Y_t)dt + \Big(a_0^2 + \varepsilon\, a_1^2 e^{\beta Y_t}\Big)^{1/2} dW_t + \int_{\mathbb{R}} z\, d\widetilde{N}_t(Y_{t-}, dz), \qquad\qquad Y_0 = y, \\ \alpha(Y_t) &= \Big(c_0 + \varepsilon\, c_1 e^{\beta Y_t}\Big) - \frac{1}{2}\Big(a_0^2 + \varepsilon\, a_1^2 e^{\beta Y_t}\Big) - \int_{\mathbb{R}} \nu(Y_{t-}, dz)\Big(e^z - 1 - z\Big), \\ \zeta &= \inf\left\{ t > 0 : \int_0^t \left(c_0 + \varepsilon\, c_1 e^{\beta Y_s}\right) ds = \mathcal{E} \right\}, \\ \nu(Y_{t-}, dz) &= \nu_0(dz) + \varepsilon\, e^{\beta Y_{t-}} \nu_1(dz) \end{aligned}$$

We call this model the "CÉV" or "Constant Lévylasticity of Variance" model, as it combines aspects of both the CEV model and exponential Lévy models. We point out some of the main features of this model below:

- For $\beta < 0$, the local volatility function $\sigma(y) = \left(a_0^2 + \varepsilon a_1^2 e^{\beta y}\right)^{1/2}$ behaves asymptotically like the CEV model $\sigma(y) \sim \sqrt{\varepsilon}\, a_1 e^{\beta y/2}$ as $y \to -\infty$ $(x \to 0)$, reflecting the fact that volatility tends to increase as the asset price X drops (the leverage effect). As $y \to \infty$ $(x \to \infty)$ the volatility function behaves like a constant $\sigma(y) \sim a_0$. This is in *contrast* to the CEV model, in which volatility function goes unrealistically close to zero as $y \to \infty$ $(x \to \infty)$.

- Jumps of size dz arrive as a Poisson process with an instantaneous state-dependent intensity of $\nu(Y_{t-}, dz)$. The state-dependent Lévy measure $\nu(y, dz) = \nu_0(dz) + \varepsilon e^{\beta y} \nu_1(dz)$ behaves asymptotically as $\nu(y, dz) \sim \varepsilon e^{\beta} \nu_1(dz)$ as $y \to -\infty$ and asymptotically like $\nu(y, dz) \sim \nu_0(dz)$ as $y \to \infty$. Thus, both the jump intensity and jump distribution can change drastically depending on the value of y.

- The process Y is killed with a state-dependent intensity of $k(Y_t) = \left(c_0 + \varepsilon c_1 e^{\beta Y_t}\right)$. For $\beta < 0$ the function $k(y)$ behaves asymptotically like $\varepsilon c_1 e^{\beta y}$ as $y \to -\infty$, reflecting the fact that a default is more likely to occur as Y drops. As $y \to \infty$ the function $k(y)$ behaves asymptotically as a constant $k(y) \sim c_0$. This particular form of killing rate was first suggested by Carr and Linetsky (2006).

- Although both the volatility $\sigma(y)$ and jump intensity $\nu(y, dz)$ are unbounded as $y \to -\infty$ (for $\beta < 0$), negative infinity is still a natural boundary for the process Y. This is due to the fact that as $y \to -\infty$ the killing rate $k(y)$ is also unbounded.

Remark 14. Note that $e_\beta \notin C_0^\infty(\mathbb{R})$. Thus, strictly speaking, the CÉV model does not belong to the class of models described in section 2. Nevertheless, for any $y_0 > -\infty$ the function e_β *is* an element of $C_0^\infty(\mathbb{R}_0)$, where $\mathbb{R}_0 := (y_0, \infty)$. Thus, we expect the results of this section to remain valid for all $y > y_0$ for ε_{y_0} small enough. The numerical experiments provided at the end of the paper support this notion.

We wish to find an expression for u_n when $\eta = e_\beta$. Noting that

$$(\psi_\mu, e_\beta \psi_\lambda) = \delta(\lambda - \mu - i\beta),$$

we see that the $(n+1)$-fold integral (3) collapses into a single integral

$$\begin{aligned} u_n &= \int_{\mathbb{R}} d\lambda \left(\sum_{k=0}^{n} \frac{e^{t\phi_{\lambda - ik\beta}}}{\prod_{j\neq k}^{n} (\phi_{\lambda-ik\beta} - \phi_{\lambda - ij\beta})} \right) \left(\prod_{k=0}^{n-1} \chi_{\lambda - ik\beta} \right) (\psi_\lambda, h)\, \psi_{\lambda - in\beta} \\ &= e_{n\beta} \int_{\mathbb{R}} d\lambda \left(\sum_{k=0}^{n} \frac{e^{t\phi_{\lambda - ik\beta}}}{\prod_{j\neq k}^{n} (\phi_{\lambda-ik\beta} - \phi_{\lambda - ij\beta})} \right) \left(\prod_{k=0}^{n-1} \chi_{\lambda - ik\beta} \right) (\psi_\lambda, h)\, \psi_\lambda . \end{aligned} \tag{9}$$

Remark 15. Although we have written the option price as an infinite series (2), from a practical standpoint, one may only compute $u^\varepsilon \approx u^{(N)} := \sum_{n=0}^{N} \varepsilon^n u_n$. For any finite N we may pass the sum through the integral appearing in (9). Thus, for the purposes of computation, the best way express the approximate option price is

$$u^\varepsilon \approx \int_{\mathbb{R}} d\lambda\, (\psi_\lambda, h)\, \psi_\lambda \sum_{n=0}^{N} \varepsilon^n\, e_{n\beta} \left(\sum_{k=0}^{n} \frac{e^{t\phi_{\lambda - ik\beta}}}{\prod_{j\neq k}^{n} (\phi_{\lambda-ik\beta} - \phi_{\lambda - ij\beta})} \right) \left(\prod_{k=0}^{n-1} \chi_{\lambda - ik\beta} \right).$$

Note, to obtain the approximate value of u^ε, *only a single integration is required.* This makes our pricing formula as efficient as other models in which option prices are expressed as a Fourier-type integral (e.g. Lévy processes, Heston model, etc.).

Numerical Results

Define the transition density $p^\varepsilon(t, y; y_0)$ and the $\mathcal{O}(\varepsilon^n)$ approximation of the transition density $p^{(n)}(t, y; y_0)$

$$p^\varepsilon(t, y; y_0) = \mathbb{E}_{y_0}\, \delta_y(Y_t), \qquad p^{(n)}(t, y; y_0) = \sum_{k=0}^{n} \varepsilon^k p_k(t, y; y_0).$$

In figures 1 and 2 we plot the approximate transition density $p^{(n)}$ for a specific model within the CÉV framework. In figure 3 we plot the implied volatility smile induced by a specific model within the CÉV framework (see equation (5) for a definition of implied volatility). The smile is generated by computing u^ε first (using Theorem 2) and then by inverting $u^{BS}(\sigma^\varepsilon)$ numerically to obtain σ^ε.

Next, for a particular model in which $\nu_0 \equiv 0$ and $c_0 = 0$ (the assumption of section 4) define the $\mathcal{O}(\varepsilon^n)$ approximation of the implied volatility

$$\sigma^{(n)} := \sum_{k=0}^{n} \varepsilon^k \sigma_k .$$

In figure 4 we provide a numerical example illustrating convergence of $\sigma^{(n)}$ to σ^ε. We compute $\sigma^{(n)}$ using Theorem 12. Implied volatility is plotted as a function of the log-moneyness to maturity ratio, LMMR := $(k - y)/t$. Convergence is fastest for values of k near y and slows as k moves away from y.

6 Conclusion

In this paper we introduce a class of local Léy models. Within our modeling framework, we obtain a formula (written as an infinite series) for the price of any European option. Additionally, we obtain an explicit expression for the implied volatility smile induced by a certain sub-class of our models. As an example of our framework, we introduce a CEV-like local Lévy model, which corrects some short-comings of the CEV model; namely (i) our choice of volatility function does not drop to zero as the value of the underlying increases and (ii) our model permits the underlying asset to experience jumps. In the CEV-like framework, we show that option prices can be computed with the same level of efficiency as other models in which option prices are computed as Fourier-type integrals.

Thanks

The author would like to thank Bjorn Birnir for his helpful comments.

References

Borodin, A. and P. Salminen (2002). *Handbook of Brownian motion: facts and formulae.* Birkhauser.

Brown, J. and R. Churchill (1996). *Complex variables and applications*, Volume 7. McGraw-Hill New York, NY.

Carr, P. and V. Linetsky (2006). A jump to default extended CEV model: An application of bessel processes. *Finance and Stochastics 10*(3), 303–330.

Chernoff, P. R. (1972). Perturbations of dissipative operators with relative bound one. *Proceedings of the American Mathematical Society 33*(1).

Cox, J. (1975). Notes on option pricing I: Constant elasticity of diffusions. *Unpublished draft, Stanford University*. A revised version of the paper was published by the Journal of Portfolio Management in 1996.

Ethier, S. and T. Kurtz (1986). Markov processes. characterization and convergence.

Friedman, B. (1956). *Principles and techniques of applied mathematics*, Volume 280. Wiley New York.

Hanson, G. and A. Yakovlev (2002). *Operator theory for electromagnetics: an introduction.* Springer Verlag.

Hoh, W. (1998). Pseudo differential operators generating markov processes. *Habilitations-schrift, Universität Bielefeld*.

Lewis, A. (2001). A simple option formula for general jump-diffusion and other exponential Lévy processes.

Lipton, A. (2002). The vol smile problem. *Risk* (February), 61–65.

Lorig, M. (2012). The exact implied volatility smile for exponential Lévy models. *Arxiv preprint arXiv:1207.0233v1*.

Mendoza-Arriaga, R., P. Carr, and V. Linetsky (2010). Time-changed markov processes in unified credit-equity modeling. *Mathematical Finance 20*, 527–569.

Reed, M. and B. Simon (1980). *Methods of modern mathematical physics. Volume I: Functional Analysis.* Academic press.

Roach, G. (1982). *Green's functions.* Cambridge Univ Pr.

Rudin, W. (1973). *Functional analysis.* McGraw-Hill, New York.

A Spectral theory of normal operators in a Hilbert space

In this appendix we summarize the theory of normal operators acting on a Hilbert space. A detailed exposition on this topic (including proofs) can be found in Reed and Simon (1980) and Rudin (1973).

Let $\mathcal{H}$ be a Hilbert space with inner product $(\cdot,\cdot)$. A *linear operator* is a pair $(\mathrm{dom}(\mathcal{A}),\mathcal{A})$ where $\mathrm{dom}(\mathcal{A})$ is a linear subset of $\mathcal{H}$ and $\mathcal{A}$ is a linear map $\mathcal{A} : \mathrm{dom}(\mathcal{A}) \to \mathcal{H}$. The *adjoint* of an operator $\mathcal{A}$ is an operator $\mathcal{A}^*$ such that $(\mathcal{A}f, g) = (f, \mathcal{A}^* g), \forall\, f \in \mathrm{dom}(\mathcal{A}), g \in \mathrm{dom}(\mathcal{A}^*)$, where

$$\mathrm{dom}(\mathcal{A}^*) := \{g \in \mathcal{H} : \exists\, h \in \mathcal{H} \text{ such that } (\mathcal{A}f, g) = (f, h)\ \forall\, f \in \mathrm{dom}(\mathcal{A})\}.$$

An operator $(\mathrm{dom}(\mathcal{A}),\mathcal{A})$ is said to be *self-adjoint* in $\mathcal{H}$ if

$$\mathrm{dom}(\mathcal{A}) = \mathrm{dom}(\mathcal{A}^*), \qquad (\mathcal{A}f, g) = (f, \mathcal{A}g) \qquad \forall\, f, g \in \mathrm{dom}(\mathcal{A}).$$

Throughout this appendix, for any self-adjoint operator $\mathcal{A}$, we will assume that $\mathrm{dom}(\mathcal{A})$ is a dense subset of $\mathcal{H}$. A densely defined self-adjoint operator is closed (see Rudin (1973), Theorem 13.9). An operator $(\mathrm{dom}(\mathcal{A}),\mathcal{A})$ is said to be *normal* in $\mathcal{H}$ if it is closed, densely defined and commutes with its adjoint: $\mathcal{A}^*\mathcal{A} = \mathcal{A}\mathcal{A}^*$. Clearly, every self-adjoint operator is a normal operator.

Given a linear operator $\mathcal{A}$, the *resolvent set* $\rho(\mathcal{A})$ is defined as the set of $\lambda \in \mathbb{C}$ such that the mapping $(\mathcal{A} - \mathrm{Id}\,\lambda)$ is one-to-one and $R_\lambda := (\mathcal{A} - \mathrm{Id}\,\lambda)^{-1}$ is continuous with $\mathrm{dom}(R_\lambda) = \mathcal{H}$. The operator $R_\lambda : \mathcal{H} \to \mathcal{H}$ is called the *resolvent*. The *spectrum* $\sigma(\mathcal{A})$ of an operator $\mathcal{A}$ is defined as $\sigma(\mathcal{A}) := \mathbb{C} \setminus \rho(\mathcal{A})$. We say that $\lambda \in \sigma(\mathcal{A})$ is an *eigenvalue* of $\mathcal{A}$ if there exists $\psi \in \mathrm{dom}(\mathcal{A})$ such that the *eigenvalue equation* is satisfied

$$\mathcal{A}\,\psi = \lambda\,\psi. \tag{10}$$

A function ψ that solves (10) is called an *eigenfunction* of $\mathcal{A}$ corresponding to λ. The *multiplicity* of an eigenvalue λ is the number of linearly independent eigenfunctions for which equation (10) is satisfied. The spectrum of an operator $\mathcal{A}$ can be decomposed into two disjoint sets called the *discrete* and *essential* [1] spectra: $\sigma(\mathcal{A}) = \sigma_d(\mathcal{A}) \cup \sigma_e(\mathcal{A})$. For a normal operator $\mathcal{A}$, a number $\lambda \in \mathbb{C}$ belongs to $\sigma_d(\mathcal{A})$ if and only if λ is an isolated point of $\sigma(\mathcal{A})$ and λ is an eigenvalue of finite multiplicity (see Rudin (1973), Theorem 12.29).

A *projection-valued measure* on the measure space $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ is a family of bounded linear operators $\{E(B), B \in \mathcal{B}(\mathbb{C})\}$ in $\mathcal{H}$ that satisfies:

1. $E(\emptyset) = 0$ and $E(\mathbb{C}) = \mathrm{Id}$.

2. $E(B)$ is an orthogonal projection. That is, $E^2(B) = E(B)$ and $E(B)$ is self-adjoint: $E^*(B) = E(B)$.

---

[1] The essential spectrum may be further decomposed into the *continuous* spectrum and the *residual* spectrum. It can be shown that the residual spectrum of an ordinary differential operator is empty (see Roach (1982), page 184).

3. $E(A \cap B) = E(A)E(B)$.

4. If $B = \bigcup_{i=1}^{\infty} B_i$ and $B_i \cap B_j = \emptyset$ for $i \neq j$ then $E(B) = \lim_{n\to\infty} \sum_{i=1}^{n} E(B_j)$, where the limit is in the strong operator topology.

5. For every in $f, g \in \mathcal{H}$ the set function $\mu_{f,g}(B) := (f, E(B)g)$ is a complex measure on $\mathcal{B}(\mathbb{C})$.

Theorem 16 (Spectral Representation Theorem)**.** *There is a one-to-one correspondence between normal operators $\mathcal{A}$ and projection-valued measures E on $\mathcal{H}$, the correspondence being given by*

$$\mathcal{A} = \int_{\sigma(\mathcal{A})} \lambda \, E(d\lambda).$$

If $g(\cdot)$ is a Borel function on $\mathbb{C}$ then

$$g(\mathcal{A}) = \int_{\sigma(\mathcal{A})} g(\lambda) E(d\lambda), \qquad \mathrm{dom}(g(\mathcal{A})) = \{f \in \mathcal{H} : \int_{\sigma(\mathcal{A})} |g(\lambda)|^2 \mu_{f,f}(d\lambda) < \infty\}. \tag{11}$$

Proof. See Rudin (1973) Theorems 12.21 and 13.33. $\square$

As a practical matter, if $\mathcal{A}$ is an differential operator acting on a Hilbert space $L^2(\mathbb{R}, dy)$, then the operators defined by (11) can be constructed by solving the *proper* and *improper* [2] eigenvalue problems

$$\begin{array}{llll} \text{proper:} & \mathcal{A}\,\psi_n = \phi_n\,\psi_n, & \phi_n \in \sigma_d(\mathcal{A}), & \psi_n \in \mathcal{H}, \\ \text{improper:} & \mathcal{A}\,\psi_\lambda = \phi_\lambda\,\psi_\lambda, & \phi_\lambda \in \sigma_e(\mathcal{A}), & \psi_\lambda \notin \mathcal{H}. \end{array}$$

For the improper eigenvalue problem one extends the domain of $\mathcal{A}$ to include functions all functions ψ for which $\mathcal{A}f$ makes sense and for which the following boundedness conditions are satisfied

$$\lim_{y\to\pm\infty} |\psi(y)|^2 < \infty.$$

After normalizing, the proper and improper eigenfunctions $\mathcal{A}$ satisfy the following orthogonality relations

$$(\psi_n, \psi_m) = \delta_{n,m}, \qquad (\psi_\lambda, \psi_{\lambda'}) = \delta(\lambda - \lambda'), \qquad (\psi_n, \psi_\lambda) = 0.$$

The operator $g(\mathcal{A})$ in (11) is constructed as follows (see Hanson and Yakovlev (2002), section 5.3.2)

$$g(\mathcal{A})f = \sum_{\lambda \in \sigma_d(\mathcal{A})} g(\phi_\lambda)\,(\psi_\lambda, f)\,\psi_\lambda + \int_{\sigma_e(\mathcal{A})} g(\phi_\lambda)\,(\psi_\lambda, f)\,\psi_\lambda d\lambda.$$

It is not always easy to evaluate divergent integrals of the form $(\psi_\lambda, \psi_{\lambda'})$ and verify that they are in fact delta functions $\delta(\lambda - \lambda')$. A method for directly obtaining properly normalised improper eigenfunctions can be found on page 238 of Friedman (1956).

---

[2] The term "improper" is used because the improper eigenvalues $\lambda \notin \sigma_d(\mathcal{A})$ and the improper eigenfunctions $\psi_\lambda \notin \mathcal{H}$ since $(\psi_\lambda, \psi_\lambda) = \infty$.

B Proof of Theorem 2

Our strategy is to show that $\mathcal{A}^\varepsilon = \mathcal{A}_0 + \varepsilon\,\eta\,\mathcal{A}_1$ generates a semigroup $\mathcal{P}^\varepsilon_t = \exp(t\mathcal{A}^\varepsilon)$. This will guarantee that $u^\varepsilon(t,y) = \mathcal{P}^\varepsilon_t h(y)$ is an analytic function of ε, which in turn, justifies the use of expansion (2). Throughout this section we will work on the Hilbert space $\mathcal{H} = L^2(\mathbb{R}, dy)$. We let $\mathrm{dom}(\mathcal{A}_i) = C_0^\infty(\mathbb{R})$ and we note that $C_0^\infty(\mathbb{R})$ is a dense subset of $\mathcal{H}$. Our analysis begins with a Theorem from Chernoff (1972):

Theorem 17. *Let $\mathcal{A}$ be the generator of a C_0 contraction semigroup $\mathcal{P}^0_t = \exp(t\mathcal{A})$ on a Banach space. Let $\varepsilon\,\mathcal{B}$ be a dissipative operator with a densely defined adjoint. Assume that the inequality*

$$\|\varepsilon\,\mathcal{B}u\| \leq a\,\|u\| + b\,\|\mathcal{A}u\|, \qquad \forall\, u \in \mathrm{dom}(A)$$

holds for some $a \geq 0$ and $b \leq 1$ (i.e., the operator $\varepsilon\,\mathcal{B}$ is $\mathcal{A}$-bounded with bound $b \leq 1$). Then the closure of $\mathcal{A}^\varepsilon := \mathcal{A} + \varepsilon\,\mathcal{B}$ generates a C_0 contraction semigroup $\mathcal{P}^\varepsilon_t = \exp(t\mathcal{A}^\varepsilon)$.

Remark 18. The operator $\mathcal{A}_0$ is the generator of a C_0 contraction semigroup $\mathcal{P}^0_t = \exp(t\mathcal{A}_0)$ on $\mathcal{H}$.

To show that $\varepsilon\,\eta\,\mathcal{A}_1$ is dissipative, the following Theorem will be useful:

Theorem 19. *Let $\mathcal{A}$ be a linear operator with domain* $\mathrm{dom}(\mathcal{A}) = C_0^\infty(\mathbb{R})$. *Then $\mathcal{A}$ satisfies the positive maximum principle if and only if*

$$\mathcal{A} = \frac{1}{2}a^2(y)\partial^2 + b(y)\partial + \int \nu(y,dz)\left(e^{z\partial} - 1 - \mathbb{I}_{\{z<R\}} z\partial\right) - c(y), \tag{12}$$

for some $a(x) \geq 0$, $b(x) \in \mathbb{R}$, $c(x) \geq 0$, $R \in [0,\infty]$ and $\nu(y,dz)$ satisfying

$$\int_{\mathbb{R}} \nu(y,dz)\left(1 \wedge z^2\right) < \infty.$$

Operators of the form (12) *are called* Lévy-type operators*.*

Proof. See Theorem 2.12 of Hoh (1998). □

Remark 20. An operator that satisfies the positive maximum principle is dissipative (see Ethier and Kurtz (1986), Lemma 4.2.1 on page 165).

Remark 21. The operator $\varepsilon\,\eta\,\mathcal{A}_1$ is clearly of the form (12). Therefore, $\varepsilon\,\eta\,\mathcal{A}_1$ is dissipative.

Remark 22. The adjoint of $\varepsilon\eta\mathcal{A}_1$, given by $(\varepsilon\eta\mathcal{A}_1)^* = \varepsilon\mathcal{A}_1^*\eta$, has domain $\mathrm{dom}(\varepsilon\mathcal{A}_1^*\eta) = C_0^\infty(\mathbb{R})$, and is therefore densely defined in $\mathcal{H}$.

Since $\mathcal{A}_0$ generates a C_0 semigroup and $\varepsilon\,\eta\,\mathcal{A}_1$ is dissipative and has a densely defined adjoint, we have only to show that $\varepsilon\,\eta\,\mathcal{A}_1$ is $\mathcal{A}_0$-bounded with $b \leq 1$.

Proposition 23. *Suppose* $\eta \in C_0^\infty(\mathbb{R})$ *and* ε *satisfies*

$$\varepsilon^2 \leq \inf_{\lambda \in \mathbb{R}} \frac{a^2 + b^2 |\phi_\lambda|^2}{\|\eta\|^2 \cdot |\chi_\lambda|^2},$$

for some $a \geq 0$ *and* $b \leq 1$ *(which is the condition given in Theorem 2). Then* $\varepsilon\, \eta\, \mathcal{A}_1$ *is* $\mathcal{A}_0$*-bounded.*

Proof. First, observe that for any $u \in C_0^\infty(\mathbb{R})$ we have

$$\|\mathcal{A}_0 u\|^2 = \int d\lambda\, |(\psi_\lambda, u)|^2 |\phi_\lambda|^2, \qquad \|\mathcal{A}_1 u\|^2 = \int d\lambda\, |(\psi_\lambda, u)|^2 |\chi_\lambda|^2, \qquad \|u\|^2 = \int d\lambda\, |(\psi_\lambda, u)|^2.$$

Thus,

$$\begin{aligned}
\varepsilon^2 \leq \inf_{\lambda \in \mathbb{R}} \frac{a^2 + b^2 |\phi_\lambda|^2}{\|\eta\|^2 \cdot |\chi_\lambda|^2} \qquad &\Leftrightarrow \qquad \varepsilon^2 \leq \frac{a^2 + b^2 |\phi_\lambda|^2}{\|\eta\|^2 \cdot |\chi_\lambda|^2} \quad \forall\, \lambda \\
&\Leftrightarrow \qquad 0 \leq a^2 + b^2 |\phi_\lambda|^2 - \varepsilon^2 \|\eta\|^2 \cdot |\chi_\lambda|^2 \quad \forall\, \lambda \\
&\Rightarrow \qquad 0 \leq \int d\lambda\, |(\psi_\lambda, u)|^2 \left(a^2 + b^2 |\phi_\lambda|^2 - \varepsilon^2 \|\eta\|^2 \cdot |\chi_\lambda|^2 \right) \\
&\qquad\qquad = a^2 \|u\|^2 + b^2 \|\mathcal{A}_0 u\|^2 - \varepsilon^2 \|\eta\|^2 \cdot \|\mathcal{A}_1 u\|^2 \\
&\Leftrightarrow \qquad \varepsilon^2 \|\eta\|^2 \cdot \|\mathcal{A}_1 u\|^2 \leq a^2 \|u\|^2 + b^2 \|\mathcal{A}_0 u\|^2 \\
&\Rightarrow \qquad \|\varepsilon \eta\, \mathcal{A}_1 u\|^2 \leq a^2 \|u\|^2 + b^2 \|\mathcal{A}_0 u\|^2 \\
&\Rightarrow \qquad \|\varepsilon \eta\, \mathcal{A}_1 u\| \leq a \|u\| + b \|\mathcal{A}_0 u\|.
\end{aligned}$$

where we have used $\|\eta\, \mathcal{A}_1 u\| \leq \|\eta\| \cdot \|\mathcal{A}_1 u\|$ in the second-to-last step. □

The proof of Theorem 2 is complete.



$$\nu_i(dz) = \frac{1}{\sqrt{2\pi s_i^2}} \exp\left(\frac{-z^2}{2s_i^2}\right) dz$$

Figure 1: A plot of the approximate transition density $p^{(n)}(t, y; 0)$ (solid blue) for different values of n. For comparison, we also plot $p^{(0)}$ (dashed black). Note that the density of Y_t has a fat tail to the left, which is expected since the volatility and jump intensity increase as $y \to -\infty$. The following parameters are used in these plots: $a_0 = 0.20$, $a_1 = 0.10$, $c_0 = 0.0$, $c_1 = 0.001$, $s_0 = 1.0$, $s_1 = 0.58$, $\varepsilon = 1$, $\beta = -0.35$, $t = 1.0$.



$$\nu_i(dz) = \frac{1}{\sqrt{2\pi s_i^2}} \exp\left(\frac{-(z-m_i)^2}{2s_i^2}\right) dz$$

Figure 2: For different values of n we plot of the approximate transition densities $p^{(n)}(t, y; y_0)$ for $y_0 = -0.6$ (solid red), $y_0 = 0.0)$ (dashed black) and $y_0 = 0.6)$ (dot-dashed blue). Note that as y_0 moves closer to $-\infty$ (i) the transition density becomes wider (ii) convergence of the approximate density $p^{(n)}$ to the exact density p^ε requires more terms (iii) the densities have fatter tails on the left than on the right. All three phenomena are due to the fact that volatility and jump-intensity rise as $y \to -\infty$. The following parameters are used in these plots: $a_0 = 0.20$, $a_1 = 0.10$, $c_0 = 0.0$, $c_1 = 0.001$, $s_0 = 0.15$, $m_0 = -0.10$, $s_1 = 0.15$, $m_1 = -0.10$, $\varepsilon = 1$, $\beta = -0.95$, $t = 1.0$.



$$\nu_i(dz) = \frac{1}{\sqrt{2\pi s_i^2}} \exp\left(\frac{-(z-m_i)^2}{2s_i^2}\right) dz$$

Figure 3: A plot of the implied volatility smile induced by a model with Gaussian jumps. Units on the horizontal axis are LMMR. Note that the steep skew, which is due to the fact that volatility and jump intensity increase as the value of the underlying drops. The following parameters are used in this plot: $a_0 = 0.20$, $a_1 = 0.10$, $c_0 = 0.0$, $c_1 = 0.001$, $s_0 = 0.2$, $s_1 = 0.1$, $m_0 = -0.2$, $m_1 = -0.1$, $\varepsilon = 1$, $\beta = -1.25$, $t = 0.50$.



$$\nu_0(dz) = 0, \qquad \nu_1(dz) = \frac{1}{\sqrt{2\pi s_1^2}} \exp\left(\frac{-(z-m_1)^2}{2s_i^2}\right) dz$$

Figure 4: We plot $\sigma^{(n)}$ (solid blue) and σ^{ε} (dashed black) as a function of LMMR. The following parameters are used in these plots: $a_0 = 0.25$, $a_1 = 0.10$, $c_0 = 0.00$, $c_1 = 0.001$, $s_1 = 0.15$, $m_1 = -0.10$, $\varepsilon = 1$, $\beta = -1.25$, $t = 0.55$, $y = -0.01$.